PREVENTING RADIATION TOXICITY THROUGH DIET

A WORLD BUSINESS ACADEMY EBOOK

BETH ELLEN DILUGLIO, MS, RDN, CCN, LDN JULIE M. BROWN, MA, LMHC

CONTENTS

RADIOACTIVE ELEMENTS: NUTRIEI

RADIOACTIVE ELEMENTS AND TH

TRACKING RADIATION RESOURCES AND REFERENCES ALKALIZING AND ACIDIFYING FOODS

PREVENTING RADIATION TOXICIT

RADIATION EXPOSURE NUTRITION

NUTRITION INTERVENTION STRAT

ABOUT THE AUTHORS

World Business Academy 308 East Carrillo Street Santa Barbara, CA 93101 www.worldbusiness.org www.safeenergyproject.org

Preventing Radiation Toxicity Through Diet Beth Ellen DiLuglio, MS, RDN, CCN, LDN Julie M. Brown, MA, LMHC

All articles ©2013 Beth Ellen DiLuglio, MS, RDN, CCN, LDN 2008-2013, All rights reserved, except for article on page 13 ©2013 Julie M. Brown, MA, LMHC 2008-2013, All rights reserved, and excluding the CDC's National Report on Human Exposure to Environmental Chemicals and the National Toxicology Program's Report on Carcinogens

All photographs ©2013 Isaac Hernández/IsaacHernandez.com

Book design: Isaac Hernández

NT IMPOSTERS	4
EIR EFFECT ON THE HUMAN BODY	6
	10 11
Y	13
N INTERVENTION STRATEGIES	14
EGIES REFERENCES	18
	20

RADIOACTIVE ELEMENTS: NUTRIENT IMPOSTERS

BETH ELLEN DILUGLIO, MS, RD, CCN, LD/N

Did you know that radioactive elements masquerade as nutrients and get absorbed by the body? Nutrients are indispensable to all processes that take place in the body, they are indispensable to life. We get our essential nutrients from food and water. When radioactive elements are released into the air or water, they mimic nutrients and find their way into our food supply and into our bodies.

WHAT IS A RADIOACTIVE ELEMENT?

A radioactive element is a chemically unstable form of an element that is formed during nuclear fission, a process that occurs during production of nuclear weapons and nuclear power. They become dangerous free radicals, emitting ionizing radiation powerful enough to steal electrons from surrounding molecules. These unstable elements can damage tissues, cell membranes and DNA. This damage can lead to altered hormone function; bone decay (osteoporosis); immune suppression; altered cell membrane function and cancer. Radioactive elements are obviously not meant for human ingestion! Some radioactive elements occur naturally, such as radon and uranium, and are found in the ground where they should stay! Many radioactive elements can have a "half" life of hundreds to thousands of years!

RADIOACTIVE ELEMENTS IN THE AIR, FOOD AND WATER

Radioactive steam and radioactive waste can be released into the air and enter our lungs, come down in the rain and end up in our water and food crops. If we don't ingest enough healthy essential nutrients, the body will absorb the radioactive imposters. For example, iodine-131, strontium-90, cesium-137 and plutonium-239 can be absorbed as if they were iodine, strontium/calcium, potassium and iron respectively. Cobalt-60, culfur-35 and cinc-65 will be taken up as if they were vitamin B12, sulfur and zinc. Low doses of radiation appear to be even more dangerous as they stealthily incorporate themselves into our bones and tissues. This is known as the "Petkau Effect."

BIOACCUMULATION CONCENTRATES RADIOACTIVE ELEMENTS AND TOXINS

Toxins accumulate as we go up the food chain. For example, plankton absorb toxins and radioactive elements and they are eaten by small fish. The small fish will be eaten by larger fish who get a "double dose." Then we eat the large fish and get an even more concentrated dose of toxins and radiation. The concentration of toxins and radioactivity in fish can be thousands of times greater than what can be measured in the water they swim in. Bioaccumulation leads to an overdose for everyone!

RADIOACTIVE ELEMENTS WREAK HAVOC IN OUR BODIES

These dangerous nutrient imposters are sent to do the jobs of healthy nutrients and become incorporated into our bones, brains, soft tissues and crucial hormone producing organs such as the ovaries, testes and pancreas. When surrounding tissues become radiated over time, the damage can become incapacitating. Radioactive elements that settle in the bone and radiate the bone marrow can lead to suppression of both red blood cells (anemia) and white blood cells (immune dysfunction).

Radiation can damage the DNA of the cell, altering the delicate genetic blueprints that lie at the base of our existence. When the genes for cell replication are damaged, the cell will divide nonstop and form cancer cells and ultimately tumors. The fetus and growing children are especially sensitive to radiation. Critical periods of development can be irreversibly disrupted, leading to altered functioning of the brain, nervous system, cardiovascular system, etc.

WHAT CAN WE DO?!

Purify Your Water through reverse osmosis, distillation or softening. These processes will remove radioactive elements. It will remove healthy nutrients as well so be sure to consume a diet rich in vitamins, minerals and phytonutrients (from plants)! Willard Water is an easy way to add major minerals and trace minerals back into your drinking water.

ANTIOXIDANTS PROTECT AGAINST RADIATION

Antioxidants protect us against the oxidizing effects of free radicals and may reduce the damage caused by radioactive elements. Plant based foods are naturally high in antioxidants as a protective measure for the plants themselves. Fresh fruits, vegetables, legumes, nuts, seeds, herbs and spices are especially good sources. Choose produce that has been allowed to ripen on the vine to maximize production of antioxidants and phytonutrients. Eating a plant based diet low on the food chain will also reduce our exposure to toxic chemicals, pesticides and radiation.

PROTECTIVE FOODS

Seaweed such as kelp, nori, dulce and sea vegetables are especially protective against uptake of radioactive iodine-131 as well as blocking strontium-90. Miso (fermented soybean paste) has an alkalizing effect and provides calcium, iron, B vitamins and zybicolin (helps detoxify and eliminate radioisotopes).

Healthy foods high in calcium can help block strontium-90 uptake (almonds, collard and turnip greens, blackstrap molasses, spinach and sardines and salmon with the bone in).

Foods high in potassium are also important (baked sweet and white potatoes, tomato paste, white beans, carrot juice, bananas and spinach are good examples).

Cruciferous vegetables (arugula, Bok choy, broccoli, Brussels sprouts, cabbage, cauliflower, collards, horseradish, kale, kohlrabi, mustards greens, radishes, turnips, wasabi and watercress) provide us with antioxidants, indoles and sulfur.

Supplements can be helpful in creating a nutrient rich environment in which radioactive imposters are crowded out. Taking a high quality supplement containing Vitamin C, Vitamin E Complex (tocopherols and tocotrienols), carotenoids, CoQ10 and selenium along with adequate levels of B vitamins, Vitamin D and essential minerals would be a prudent start. Be sure to discuss supplementation with a qualified healthcare practitioner.

Glutathione is a master antioxidant and is actually produced by our bodies from glutamine, glycine and N-acetylcysteine. Cordyceps, Gotu Kola, milk thistle and alpha lipoic acid have also been shown to increase glutathione levels. A topical liposomal cream containing glutathione can also boost our internal levels.

So, in our struggle to stay healthy, it's important not to be afraid, but instead Be Aware!



RADIOACTIVE ELEMENTS AND THEIR EFFECT ON THE HUMAN BODY

BETH ELLEN DILUGLIO, MS, RD, CCN, LD/N

THE AMERICAN DIETETIC ASSOCIATION'S POSITION PAPER

"It is the position of the American Dietetic Association to encourage environmentally responsible practices that conserve natural resources, minimize the quantity of waste that is generated, and have the least adverse effect on the health of all living organisms and the environment. All components of the food system, from farmer to consumer, are affected by the availability and cost of energy and the availability and quality of water. [Harmon AH, Gerald BL; American Dietetic Association. Position of the American Dietetic Association: food and nutrition professionals can implement practices to conserve natural resources and support ecological sustainability. J Am Diet Assoc. 2007 Jun;107(6):1033-43. PubMed PMID: 17571455].

BIOACCUMULATION

Toxins and contaminants concentrate as we go up the food chain from plants to animals/seafood up to humans.

Larger flesh eating fish and animals will have a higher concentration of mercury, radiation, pesticides, etc.

Synergistic effect of complexing radiation with other harmful chemicals can increase 1000 fold.

WHAT ARE RADIOACTIVE ELEMENTS?

Many elements from the Periodic Table are MINERALS that are essential to human health.

For all elements, there are varying forms known as isotopes, determined by differing numbers of neutrons.

Unstable isotopes are radioactive and release energy (electromagnetic waves) and particles in order to become stable.

Radioactive isotopes are produced from nuclear fission and can be taken up by the body as if they were their essential mineral counterparts.

For example, iodine-131, strontium-90, cesium-137 and plutonium-239 can be absorbed as if they were iodine, strontium/calcium, potassium and iron respectively.

Once in the body, they travel through the bloodstream and are incorporated into living tissue.

As radioisotopes "decay", they release ionizing radiation (a known carcinogen) that is strong enough to remove electrons from surrounding molecules or atoms.

The resulting free radical can damage cell membranes and DNA, the blueprint of the cell

Low doses of radiation are more damaging than a direct hit on the cell's DNA (which can be repaired guickly), known as the Petkau effect.

PETKAU EFFECT

In 1972 Dr. Abram Petkau discovered that biological damage to cells is greater from ionizing radiation delivered in low doses over long periods of time versus high intensity exposure over a short period of time.

The cause of the long term damage was from free radical. Many foods and antioxidants can help protect us from free radical damage. Several elements are essential to human health.

RADIOISOTOPES AND THEIR HALF-LIFE

The half-life of a radioactive substance is the amount of time it takes for half of it to decay. Half of the original amount remains and one half of that half will then decay in a "half-life", etc. Unfortunately, radioactive elements persist in the environment and in our bodies.

EFFECTS ON THE DEVELOPING FETUS

Damaging effects of radioisotopes are magnified in the developing fetus.

Dr. Alice Stewart demonstrated correlation of x-rays during pregnancy to increased incidence of leukemia in the offspring. [Stewart A. Aetiology of Childhood Malignancies. Br Med J. 1961 Feb.

18;1(5224):452-60. PubMed PMID: 20789069]

Critical Periods of development are especially vulnerable to irreversible damage.

CRITICAL PERIODS OF DEVELOPMENT

Usually a period of rapid cell development, Critical Periods are particularly vulnerable to disruptive influences that can have permanent effects

http://thyroidbook.com/blog/radiation-nutritional-intervention/

http://drknews.com/

TRACKING RADIATION

lonizing radiation is a known human carcinogen. The "St. Louis Baby Teeth Survey" (conducted in the 1950's and 1960's) demonstrated that radioactive fallout from nuclear bomb testing in the USA was indeed being absorbed by humans.

The presence of radioactive Strontium-90 in deciduous teeth was an indicator/marker of how much radiation the mother was exposed to during pregnancy. This study and scientific debate on the issue (Dr. Ernest Sternglass - June 1963 issue of Science) led to a partial test ban treaty on nuclear bomb testing between the USA and the USSR despite the ongoing "Cold War."

THE "ST. LOUIS BABY TEETH SURVEY"

Tracking radioactive elements from nuclear fission was first conducted between the 1940's and 1960's during the nuclear bomb test years. Strontium-90 (Sr-90) was chosen as a marker because it could be tracked in human teeth and bones. Sr-90 is the radioactive isotope of Strontium (a nutrient similar to calcium in it's roles in the body and the bone).

The "St. Louis Baby Teeth Survey" demonstrat-Partial Test Ban Treaty. ed an increase in Sr-90 in the teeth of children born during nuclear weapons testing in the As Professor Emeritus of Radiological Physics at 1950's and 60's. The study collected 385,000 the University of Pittsburgh Medical School, Dr. baby teeth and completed analysis on 60,000 Sternglass has written numerous articles on the of them revealing that the average Sr-90 for health effects of low-level radiation. children born in 1964 was 50 times greater than His 1981 book Secret Fallout: Low-level Radiafor those born in 1951, the first birth year for tion from Hiroshima to Three Mile Island estabwhich teeth were measured. lished him as a pioneer in the study of the health Miraculously, 85,000 of the teeth, collected but effects of low-level radiation.

not analyzed, were discovered in St. Louis this **RPHP RADIATION.ORG** past spring. Washington University, involved with Five books published by RPHP research associates summarize hundreds of articles in peer-reviewed journals dealing with these impacts of ingested, low-level fission products--products which did not exist in nature prior to the Nuclear Age. In addition to the effects upon the immune response THE RPHP TOOTH FAIRY PROJECT of all age groups, the very young have been especially affected. RPHP has repeatedly pointed out Dr. J. Sternglass, PhD (cofounder and chief the radiation-induced damage apparent in official vital statistics, tracing changes in infant mortality rates and underweight live births in the postwar period, especially during the aboveground nuclear test years of the 1950s and the 1960s.

the initial St. Louis study, has sent the teeth to the Radiation and Public Health Project (RPHP) Tooth Fairy Project where they can be used in a prospective study of the health effects of low level radiation and Sr-90 ingestion. technical officer of the RPHP) and colleagues conducted a second "Baby Teeth Study- The Tooth Fairy Project" 1997 - present. Data indicate that levels of Sr-90 in **baby teeth** are as high today in counties close to nuclear power plants as they

were during above ground nuclear bomb testing. Sr-90 in drinking water was also significantly higher in these counties.

REVERSE OSMOSIS FOUND TO REMOVE SR-90

Sr-90 is mistaken for calcium and settles in the bone, radiating bone marrow, damaging red blood cells, white blood cells... contributing to childhood cancer, immune deficiency and osteoporosis.

www.radiation.org

CANCER: RELATION OF PRENATAL RADIATION TO DEVELOPMENT OF THE **DISEASE IN CHILDHOOD**

STERNGLASS EJ. CANCER: RELATION OF PRENATAL RADIATION TO DEVELOPMENT OF THE DISEASE IN CHILDHOOD. SCIENCE. 1963 JUN 7;140(3571):1102-4. PUBMED PMID: 13983978.

In 1963, Dr. Sternglass was invited to testify before the congressional Joint Committee on Atomic Energy, as to how the exponential increase in strontium-90 in baby teeth caused by bomb-test fallout was associated with increased childhood leukemia. His research and testimony played a role in President Kennedy's decision to sign the

RPHP has also been able to track the radiation-induced damage done to the hormonal and immune systems of the 80 million baby boomers born between 1945 and 1965 in each of the post war decades, revealing the various epidemiological anomalies: In the 1950s, children born after the enormous initial exposure to nuclear fission products began to experience epidemic increases in childhood cancer in the ages 5 to 9.

A total of 422 nuclear weapons were detonated in the atmosphere by the United States (206 tests) and the Soviet Union (216 tests) before largescale testing ended with the 1963 Partial Test Ban Treaty. Yield from the six largest Soviet tests alone totaled 136.9 megatons, or the equivalent of nearly 4,000 Hiroshima and Nagasaki bombs (36 kilotons). As testing escalated, deposition of fallout in human bodies was documented around the world. Risks to health from large-scale atmospheric nuclear weapons testing are still relatively unknown.

HEALTH STATISTICS EMERGE DECADES LATER

On October 20, 2009, RPHP released its report on the cancer risk from atomic bomb testing: CANCER RISK TO AMERICANS FROM ATOMIC TEST FALLOUT, A CASE CONTROL STUDY OF STRONTIUM-90 IN BABY TEETH. Our new study finds high levels of radioactive Strontium-90 in baby teeth of U.S. "Baby Boomers" who have died of cancer. The study marks the first attempt to estimate cancer risk from bomb fallout by measuring actual levels in bodies of Americans.

ELEVATED IN VIVO STRONTIUM-90 FROM NUCLEAR WEAPONS TEST FALLOUT AMONG CANCER DESCENDANTS: A CASE CONTROL STUDY OF DECIDIOUS TEETH

A sample of 85,000 deciduous teeth collected from Americans born during the bomb-testing years assessed risk by in vivo measurement of residual strontium-90 (Sr-90) concentrations, using liquid scintillation spectrometry. The authors' analysis included 97 deciduous teeth from persons born between 1959 and 1961 who were diagnosed with cancer, and 194 teeth of matched controls. Average Sr-90 in teeth of persons who died of cancer was significantly greater than for controls (OR = 2.22; p < 0.04). This discovery suggests that many thousands have died or will

die of cancer due to exposure to fallout, far more than previously believed.[Mangano JJ, Sherman JD. Elevated in vivo strontium-90 from nuclear weapons test fallout among cancer decedents: a case-control study of deciduous teeth. Int J Health Serv. 2011;41(1):137-58. PubMed PMID: 213197261

FOOD IRRADIATION THREATENS PUBLIC HEALTH, NATIONAL SECURITY

...As well documented since the 1960s, these massive doses of ionizing radiation produce profound chemical changes in meat. These include elevated levels of the carcinogenic chemical **benzene**, and also the production of unique new chemicals, known as radiolytic products, some of which have been implicated as carcinogenic.

Additionally, irradiated food has been shown to induce genetic damage in a wide range of studies ... Furthermore, as admitted by USDA's Aqricultural Research Service, irradiation results in major losses of vitamins, particularly A, C, E and the B complex. (Epstein SS, Hauter W. Preventing pathogenic food poisoning: sanitation, not irradiation. Int J Health Serv. 2001;31(1):187-92. PubMed PMID: 11271643).

THE CENTER FOR DISEASE CONTROL AND PREVENTION'S REPORT ON HUMAN EXPOSURE TO ENVIRONMENTAL CHEMICALS

The first Report (March 2001) presented exposure data for 27 chemicals from NHANES 1999; this Second Report (January 2003) presents human exposure data for 116 chemicals (including the 27 in the first Report) from NHANES 1999-2000. The Second Report also presents exposure data for the U.S. population divided into age, gender, and race/ethnicity groups. Radioactive elements and toxins have a synergistic effect and are more deadly when combined.

The first National Report measured lead, mercury, cadmium, and other metals; dialkyl phosphate metabolites of organo-phosphate pesticides; cotinine; and phthalates. The Second Report includes these chemicals and adds:

Polycyclic aromatic hydrocarbons (PAHs)

Dioxins, furans, and coplanar

Polychlorinated biphenyls (PCBs)

Non-coplanar PCBs Phytoestrogens Herbicides Selected organophosphate pesticides Organochlorine pesticides Carbamate pesticides

Pest repellents and disinfectants

The Fourth National Report on Human Exposure to Environmental Chemicals 2009 and the Updated Tables, February 2011, together are the most comprehensive assessment of environmental chemical exposure in the U.S. population. Since 1999, CDC has measured 219 chemicals in people's blood or urine. The Fourth Report, 2009, includes the findings from national samples for 1999-2000, 2001-2002, and 2003-2004.

http://www.cdc.gov/exposurereport/

• Neutrons were first listed in the Eleventh Re-Agency for Toxic Substances and Disease Regport on Carcinogens (2004). istry, 4770 Buford Hwy NE, Atlanta, GA 30341

Contact CDC: 800-232-4636 / TTY: 888-232-6348

Ionizing Radiation

CAS ID #: HZ1800-45-T

Affected Organ Systems: Dermal (Skin), Developmental (effects during periods when organs are developing), Musculoskeletal (Muscles and Skeleton), Neurological (Nervous System), Renal (Urinary System or Kidneys), Respiratory (From the Nose to the Lungs)

Cancer Effects: None

10 Jun 2011: News Release. New Substances Chemical Classification: Radionuclides (radio-Added to HHS Report on Carcinogens - The U.S. active materials) Department of Health and Human Services today Summary: Ionizing radiation is any one of sevadded eight substances to its Report on Carcinogens, a science-based document that identifies chemicals and biological agents that may put people at increased risk for cancer.

eral types of particles and rays given off by radioactive material, high-voltage equipment, nuclear reactions, and stars. The types that are normally important to your health are alpha particles, beta "Reducing exposure to cancer-causing agents particles, x rays, and gamma rays. Alpha and beta is something we all want, and the Report on particles are small, fast-moving bits of atoms that Carcinogens provides important information on a radioactive atom gives off when it changes into substances that pose a cancer risk," said Linda another substance. X rays and gamma rays are Birnbaum, Ph.D., director of both the National Intypes of electromagnetic radiation. These radistitute of Environmental Health Sciences (NIEHS) ation particles and rays carry enough energy to and the National Toxicology Program (NTP). "The knock out electrons from atoms and molecules NTP is pleased to be able to compile this report." (such as water, protein, and DNA) that they hit or

pass near. This process is called ionization, which is why this radiation is called "ionizing radiation."

IONIZING RADIATION IS A KNOWN HUMAN CARCINOGEN

lonizing radiation is electromagnetic radiation that has sufficient energy to remove electrons from atoms. Ionization results in the production of negatively charged free electrons and positively charged ionized atoms. Ionizing radiation can be classified into two categories: photons (X-radiation and gamma radiation) and particles (alpha and beta particles and neutrons). Five types or sources of ionizing radiation are listed in the Report on Carcinogens as known to be human carcinogens, in four separate listings:

• X-radiation and gamma radiation (included in one listing) were first listed in the Eleventh Report on Carcinogens (2004).

• Radon and its isotopic forms radon-220 and radon-222, which emit primarily alpha particles, were first listed in the Seventh Annual Report on Carcinogens (1994).

• Thorium dioxide, which decays by emission of alpha particles, was first listed in the Second Annual Report on Carcinogens (1981).

NATIONAL TOXICOLOGY PROGRAM'S **REPORT ON CARCINOGENS 12TH EDITION**

FOR MORE INFORMATION CONTACT: ROBIN MACKAR, NIEHS (919) 541-0073

John Bucher, Ph.D., associate director of the NTP added, "This report underscores the critical connection between our nation's health and what's in our environment."

Read more about how ionizing radiation damages cells, tissues, and organs. http://ntp.niehs. nih.gov/ntp/roc/twelfth/profiles/lonizingRadiation.pdf. Accessed September 27, 2013.

RESOURCES AND REFERENCES

1. Morales A, Miranda M, Sánchez-Reyes A, Biete A, Fernández-Checa JC. Oxidative damage of mitochondrial and nuclear DNA induced by ionizing radiation in human hepatoblastoma cells. Int J Radiat Oncol Biol Phys. 1998 Aug 1;42(1):191-203.

2. Morse ML, Dahl RH. Cellular glutathione is a key to the oxygen effect in radiation damage. Nature. 1978 Feb 16;271(5646):660-2.

3. Koval' TV, Nazarova OO, Matyshevs'ka OP. [Change in glutathione content in rat thymocytes under apoptosis induced by H2O2 or irradiation]. Ukr Biokhim Zh. 2008 Mar-Apr;80(2):114-9. Ukrainian.

4. Manu KA, Leyon PV, Kuttan G. Studies on the protective effects of Boerhaavia diffusa L. against gamma radiation induced damage in mice. Integr Cancer Ther. 2007 Dec;6(4):381-8.

5. Verma P, Sharma P, Parmar J, Sharma P, Agrawal A, Goyal PK. Amelioration of Radiation-Induced Hematological and Biochemical Alterations in Swiss Albino Mice by Panax ginseng Extract. Integr Cancer Ther. 2010 Aug 11.

6. Lee TK, O'Brien KF, Wang W, Johnke RM, Sheng C, Benhabib SM, Wang T, Allison RR. Radioprotective effect of American ginseng on human lymphocytes at 90 minutes postirradiation: a study of 40 cases. J Altern Complement Med. 2010 May;16(5):561-7.

7. Park E, Hwang I, Song JY, Jee Y. Acidic polysaccharide of Panax ginseng as a defense against small intestinal damage by whole-body gamma irradiation of mice. Acta Histochem. 2011 Jan;113(1):19-23.

8. Goel HC, Bala M, Prasad J, Singh S, Agrawala PK, Swahney RC. Radioprotection by Rhodiola imbricata in mice against whole-body lethal irradiation. J Med Food. 2006 Summer;9(2):154-60.

9. Arora R, Chawla R, Sagar R, Prasad J, Singh S, Kumar R, Sharma A, Singh S, Sharma RK. Evaluation of radioprotective activities Rhodiola imbricata Edgew-a high altitude plant. Mol Cell Biochem. 2005 May;273(1-2):209-23.

10. Kuttan G. Use of Withania somnifera Dunal as an adjuvant during radiation therapy. Indian J Exp Biol. 1996 Sep;34(9):854-6.

WEBSITES

Radiation.org Uspirg.org Ucsusa.org Cehn.org EWG.org Ourstolenfuture.org

EPA Health Effects of Radiation http://www.epa. gov/radiation/understand/health effects.html

Pubmed Collection: More Radiation and Nutrition Resources: http://www.ncbi.nlm.nih.gov/ sites/myncbi/collections/public/10Mdxrf0gg-L7bPluBqVKZjv5d/

BOOKS

Shannon, Sara. Diet for the Atomic Age : How to Protect Yourself from Low-Level Radiation (Revised Edition). New York: Instant Improvement, Inc. 1993.

Shannon, Sara. Radiation Protective Foods: A Menu For The Nuclear Age. Bloomington: AuthorHouse. 2012.

Cline, John C. and DiLuglio, Beth Ellen. Electromagnetic Hypersensitivity and Implications for Metabolism. In: Kohlstadt, I., ed. Advancing Medicine with Food and Nutrients, 2nd edition. Boca Raton, FL: CRC Press, pp. 799-820.

Whitney, Ellie and Sharon R. Rolfes. Understanding Nutrition. 11th edition. USA: Thomson, Wadsworth, 2008.

Levine, Buck, Ph.D., R. D. Environmental Nutrition. Vashon Island, Washington: Hingepin. 1999.

ALKALIZING AND ACIDIFYING FOODS

MOST ALKALIZING Baking soda Sea salt Mineral water Umeboshi plum Pumpkin seed Lentils Brocoflower Seaweed Sea vegetables Onion Miso Daikon Taro Root Burdock Lotus root Sweet potato/Yam Lime Nectarine Persimmon Raspberry Watermelon Tangerine Pineapple

> Prepared by Dr. Russell Jaffe, MD, PhD, used with permission. Complete tables, updates and reprints available from ELISA/ACT Biotechnologies. 109 Carpenter Drive, Suite 100, Sterling, VA 20164. 1-800-553-5472

MOST ACIDIFYING Table Salt Yeast/hops/malt Beer Sugar Cocoa White/acetic vinegar Processed cheese Ice cream Beef Lobster Pheasant Barley Processed flour Cottonseed oil Hazelnut Walnut Brazil nut Fried foods Soybean Carob



PREVENTING RADIATION TOXICITY

JULIE M. BROWN, MA, LMHC

Water: drink reverse osmosis water to protect yourselves and your families from radiation.
Molecules of radiation are extremely small and are only screened out by reverse osmosis. You can add ¼ tsp of sea salt into a gallon of R.O. water in order to add minerals back into your water, but if you are eating sea vegetables, you will be getting sufficient minerals.
Hydrate and Anti-Oxidant Extreme.
Eating cruciferous vegetables, broccoli, cauliflower, kale and brussel sprouts increase antioxidant and detoxification support.
Eat a high fiber diet, high pectin foods like carrots, sunflower seeds and apples take toxins, including radioactive particles out of your body.

Eating a plant-based diet, anything that grows To protect yourself from lodine-131 expoout of the ground, is very important as well. sure, take detoxified iodine (iodinesource. com) or you can take 5 kelp tablets a day. Radioactive pollutants in the environment Seaweed, plentiful in Iodine 127, protects the become more concentrated in foods like milk, cheese and meat. This is one reason why a livebody from absorbing lodine 131. It has been recorded that 20 years after the Chernobyl difood, plant-based diet, eating low on the food saster, the Polish people treated with lodine chain, is the safest and best diet as this point in history and the foreseeable future. 127 have almost 1000 times less thyroid disease than neighboring countries even further than Poland from the disaster site.

To protect yourself from cesium poisoning, consume plenty of high potassium foods, as potassium competively inhibits cesium uptake. Foods high in potassium include sea vegetables, avocados, leafy green vegetables, bananas, or make yourself a potassium tonic, juice of 3 carrots, ½ bunch spinach, ½ bunch parsley and 3 stalks celery.

To protect yourself from plutonium poisoning, eat lots of dulse and sea algaes such as spirulina, and chlorella. Miso soup has shown to be protective against plutonium. NCD, which comes from the mineral, zeolite, at 10 drops 4 times daily, is excellent for taking most radioactive elements out of the body.

Eating foods high in calcium and/or taking a high quality vit/min formula will protect the body from Strontium-90 uptake. Some foods high in calcium are spinach, almonds figs and canned Alaskan salmon, bone-in. Sr-90 acts like calcium and enters through the bone into the bone marrow. Drinking reverse osmosis water and having sufficient calcium in your bones will protect you from Sr-90.

Eat foods high in antioxidants and/or take antioxidant supplements, with vitamins C, E, beta carotenes, selenium, zinc, CoQ10, silymarin, N-acetyl-cysteine, and curcumin, and green and white tea. Two products recommended by Dr. Gabriel Cousens are MegaSources: Dr. Russell Blaylock and Dr. Gabriel Cousens.



13

RADIATION EXPOSURE NUTRITION INTERVENTION STRATEGIES

BETH ELLEN DILUGLIO, MS, RD, CCN, LD/N (NUTRITION MISSION™)

PROTECTIVE FACTORS IN RADIATION EXPOSURE

Well-nourished individuals are best equipped to block uptake, excrete radionuclides, and repair DNA damage from EMR exposure. Nutrient deficiency can be detrimental while nutrient sufficiency and saturation have protective effects.

Alkalizing Diet	Goal: arterial blood pH 7.45, first morning urine pH 6.7-7.5. Fruits and vegetables in general tend to be the most alkalizing foods, while sug- ar, meat, dairy, fried foods, and trans-fats are most acid-forming [1]. Mineral sufficiency is also crucial.
Antioxidants and Phytonutri- ents	Antioxidants, vitamins C and E, alphalipoic acid, ubiquinol, superox- ide dismutases, glutathione [2].
Detoxification Support	 Phase I: B-complex, glutathione, branched-chain amino acids, flavo- noids, phospholipids. Phase II: glycine, taurine, glutamine, N-acetylcysteine, cysteine, me- thionine, methyl donors. Intermediary: Vitamins C and E, selenium, copper, zinc, manganese, CoQ10, thiols, bioflavonoids, silymarin, pycnogenol.
Fiber	Insoluble (cellulose, lignin) and soluble (pectins, gums, gels) fiber plays an important role in radioprotection. Fiber adds bulk, speeds gastrointestinal transit time, absorbs toxicants, and promotes the growth of protective, probiotic bacteria.
Herbs and Spices	Herbs and spices are rich sources of antioxidants and phytonutri- ents that can inhibit carcinogen formation and activation, upregulate phase II detoxification enzymes, inhibit oxidation and inflammation, and demonstrate anti-tumor activity [3]. Herbs and spices studied for their protective antioxidant and anti-inflammatory effects include gar- lic, chives, onions, parsley, sage, rosemary, thyme, watercress, horse- radish, dill, bay leaves, turmeric, and tea.
Legumes	Legumes (dried beans) contain minerals, chelating-phytates and radi- oprotective protease inhibitors [4, 5].
Miso	Miso, a lactobacillus-fermented paste made from soybean and sea salt (aged ~18 months), has an alkalizing effect and is a source of cal- cium, iron, B vitamins, and zybicolin which helps bind and eliminate radionuclides [4, 5].
Nuts and Seeds	Nuts and seeds provide full spectrum vitamin E, B-complex, calcium, magnesium, potassium, iron, zinc, fiber, pectin, phytates, and ome- ga-3 fatty acids. Sesamol from sesame seeds was also found to be radioprotective a`nd exhibited a free-radical scavenging capacity 20 times that of melatonin [6].
Selective Uptake	Stable elements will block uptake of radionuclides: Calcium blocks Sr-90; Cobalamin blocks cobalt-60; Iodine blocks iodine-131; Iron blocks plutonium 238,239; Potassium blocks cesium-137; Sulfur blocks sulfur-35; Zinc blocks zinc-65.

Tempeh	Tempeh, a ferm tates, and analo
Vegetables	Vegetables con The Brassicacea cauliflower, Bru which protect c
Vegetables, Sea	Sea vegetables, and nori from ur to reduce intest
Water Purification	Reverse osmosi clides.
Whole Grains	Whole grains, a tates (which bin

PROTECTIVE SUPPLEMENTS IN RADIATION EXPOSURE

Adaptogens	Adaptogens (a zandra, rhodio erhaavia diffus docrine-immur
Aged Garlic Extract	Aged Garlic E enges reactive zymes and cell age, and inhibi
Alpha-lipoic Acid	Alpha-lipoic ac and E, increase lar and extrace ple exposed to at OSU recomr
Antioxidant Enzymes	Radiation dep glutathione p dismutases (SG als copper, zin mutase (MnSG SOD) are key produced enc ine, is also ava glutathione). (against ionizin

nented soy product, contains beneficial bacteria, phyogues of B12 that can block cobalt-58,60.

ntain fiber, minerals, phytonutrients, and antioxidants. eae family (broccoli, cabbage, collard, kale, watercress, ussels sprouts, radish, etc.) contains sulfur compounds cells from radiation.

s, (including agar, dulse, hijiki, irish moss, kelp, wakame, uncontaminated sources) are rich in minerals and found stinal absorption of Sr-90.

sis, distillation, and ion exchange can remove radionu-

as tolerated, provide vitamins, minerals, fiber, and phynd radionuclides but can also bind nutritive minerals).

astragalus, ashwagandha, ginseng, eleutheroccus, schiola, maitake and reishi mushrooms, holy basil, and bosa) exert radioprotective effects and modulate neuroenne communication.

Extract (AGE) protects against ionizing radiation, scave oxygen species, enhances cellular antioxidant enllular glutathione, protects DNA from free-radical dambits multi-step carcinogenesis [7].

cid (ALA), a potent antioxidant, regenerates vitamins C es intracellular glutathione, and protects the intracelluellular environment [8]. "ALA may be beneficial to peoo high levels of radiation" [9]. The Linus Pauling Institute mends 200-400 mg/d for healthy people.

pletes antioxidants and antioxidant enzymes such as peroxidase and glutathione reductase, superoxide SODs), and catalase. SODs utilize the essential minernc, manganese, and iron. Manganese superoxide dis-SOD) and copper-zinc superoxide dismutase (CuZny intracellular antioxidants. Glutathione, a tri-peptide adogenously from glutamic acid, glycine, and cystevailable in IV, topical, and oral form (as stable s-acetyl Glutathione and MnSOD are particularly protective ing radiation [10, 11].

Ascorbic Acid	Ascorbic acid (vitamin C) is a primary antioxidant and regenerates other antioxidants. Radiation and heavy metal exposure, stress, in- fection, and temperature changes increase requirements. The Linus Pauling Institute at OSU recommends a base dose of 250 mg vita-	Potasium Iodide	Potassium iodide active iodine. FD drugs/emergen ucm072265.htm
Astaxanthin	min C BID. For optimal health, Dr. Pauling recommends 2.3 grams or more per 2,500 Kcals [12].Astaxanthin is a xanthophyll carotenoid primarily found in marine or- ganisms such as microalgae (Haematococcus pluvialis, and Chlorella zofingiensis) krill, trout, salmon, shrimp, crayfish, and crustaceans, as	Seaweed, Sodium Alginate	Supplementing etables was fou blocks intestina tium, and incre- um metabolism
Beta-glucans	 well as bee propolis [13]. Astaxanthin possesses radioprotective, antioxidant, and immune-stimulating effects [14]. Beta-glucans are plant and microbe-based polysaccharides found in barley, oats, baker's yeast, and mushrooms. Beta-glucans stimulate hematopoiesis following ionizing radiation [15], stimulate immune cells, and down-regulate immunosuppressive cells [16]. Administration prior to, and within 24 hours of radiation exposure reduced signs of radiation sickness, enhanced immune cell response [17, 18], and may be considered for use during nuclear emergencies and RT [19,20]. 	Spirulina	Spirulina plant algae [38], was ar melt-down in a dose of 5 g p cocyanin conte effects, comple facilitating thei exerts dose-de cola recommen 10-20 g/d for t
Chlorella	Chlorella species are a type of single-celled fresh water green algae known to bind and eliminate toxins and heavy metals [21]. Chlorella's radioprotective, bioprotective, and antioxidant effects have been doc- umented in several studies [22-27]. Chlorella should be consumed in "broken cell wall" form to enhance its bioavailability. Dr. Joseph Mer- cola recommends at least 4 g daily (from uncontaminated sources) combined with fresh cilantro for a synergistic effect.	Vitamin D (1,25-dihydroxy-vitamin D3)	n D Vitamin D, a ho
Fatty Acids	Conditionally-essential omega-3 fatty acids EPA and DHA are consid- ered anti-inflammatory and immune-supportive with EPA specifical- ly protective against UV radiation [28, 29]. Cold water, oily fish such as mackerel, sardines, salmon, and purified fish oils are excellent source of EPA and DHA, while flaxseed, chia seed, hemp seed, and English walnuts are excellent sources of their precursor – alpha-lino- lenic acid. Flaxseeds were found to mitigate the negative effects of radiation, including inflammation, pulmonary fibrosis, and cytokine secretion [30]. USDA "Adequate Intake" of omega-3 fatty acids is 1.1-1.6 g/d for adults. Eating omega-3 rich seafood or consuming 2 g of high-quality fish oil is recommended several times per week by the Linus Pauling Institute at OSU.		cological agent tion damage ar [43]. Endogend exposure, amore exposed, and u less than 20 ng ng/mL. "Studies 1,100 to 4,000 tion [25(OH)D] cer risk" [44] wh ed to achieve a cohort. Few for be indicated. T
Genistein	Genistein, a phytonutrient found in soybeans, exerts radioprotec- tive, antioxidant, and anti-tumor effects [31]. Genistein applied fol- lowing radiation was found to mitigate oxidative damage, lung fi- brosis, and pneumonitis [32].	Zeolites	supplement wir serum level of a Zeolites, hydrate
Melatonin	Melatonin, produced primarily in the pineal gland from serotonin, possesses radioprotective and antioxidant properties in addition to its role in circadian rhythm regulation [33, 34]. Recommended doses range from 0.5-6 mg at bedtime [35].		occur naturally b ion-exchange ag www.chemistrye have been utiliz tissues as well ag

ide protects the thyroid during acute exposure to radio-FDA guidelines must be followed. [http://www.fda.gov/ encypreparedness/bioterrorismanddrugpreparedness/ tm]

ng with sodium alginate from kelp and other sea vegound to have a profound radioprotective effect as it nal absorption and bone uptake of radioactive stronreases Sr-90 excretion without interfering with calcim [36, 37].

ntensis, a radioprotective, unicellular blue-green as used extensively following the Chernobyl nuclein workers and children with radiation sickness at per day for 45 consecutive days [39-41]. The phyitent of Spirulina contributes to its radioprotective plexing with heavy metals and radionuclides and eir excretion. Spirulina inactivates superoxide and dependent anti-inflammatory effects [42]. Dr. Merends preventative doses of 3 g/d and increase to r therapeutic purposes [39].

normone produced in the body from cholesterol in of UV light, can be administered in supplement form ividuals from background radiation as well as nucle-Protective mechanisms include "cellular differentiamunication, Programmed Cell Death (PCD) (apopophagy) and antiangiogenesis... vitamin D... should d among the prime (if not the primary) nonpharmants that offer protection against sublethal low radiaand, in particular, against radiation-induced cancer" nous synthesis is inhibited by inadequate sunlight ount of body fat, skin pigmentation, amount of skin use of sun block. Deficiency occurs at a serum level ng/mL and sufficiency occurs in the range of 33-80 es indicate that intake of vitamin D in the range from 0 IU/d and a serum 25-hydroxyvitamin D concentra-)] from 60-80 ng/mL may be needed to reduce canwhile a supplemental dose of 9,600 IU/d was needat least 40 ng/mL in 97.5% of a community-based oods contain vitamin D and supplementation may The Linus Pauling Institute recommends that adults vith at least 2,000 IU (50 mcg) daily and maintain a f at least 80 nmol/L (32 ng/mL).

ated aluminum silicates with cation exchange capacity, y but also can be synthesized and are frequently used as agents, filters, and water softeners. HYPERLINK "http:// yexplained.com" Both natural and synthetic zeolites ilized in the removal of radionuclides from biological as from water supply systems [45, 46].

NUTRITION INTERVENTION STRATEGIES REFERENCES

1. Jaffe, RM. http://www.elisaact.com/pdfs/EAB_AlkalineWay.pdf. Accessed November 25, 2011.

2. American Institute for Cancer Research. www.aicr.org. Accessed November 13, 2011.

3. Craig WJ. Health-promoting properties of common herbs. Am J Clin Nutr. 1999 Sep;70(3 Suppl):491S-499S. Review. PMID: 10479221.

4. Shannon S. Diet for the Atomic Age. Instant Improvement, Inc. 1987.

5. Shannon, S. Radio-protective Foods- A Guide to Designing a Radiation Shield Menu of Healthy Eating. Authorhouse. 2012.

6. Mishra K, Srivastava PS, Chaudhury NK. Radiat Res. 2011 Nov;176(5):613-23. Sesamol as a potential radioprotective agent: in vitro studies. PMID: 21899433.

7. Borek C. Antioxidant health effects of aged garlic extract. J Nutr. 2001 Mar;131(3s):1010S-5S. PMID: 11238807.

8. Packer L, Tritschler HJ, Wessel K. Neuroprotection by the metabolic antioxidant alpha-lipoic acid. Free Radic Biol Med. 1997;22(1-2):359-78. Review. PMID: 8958163.

9. Natural Standard Database. Alpha Lipoic Acid. http://www.naturalstandard.com/databases/herbssupplements/alphalipoicacid.asp?. Accessed November 6, 2011.

10. Morse ML, Dahl RH. Cellular glutathione is a key to the oxygen effect in radiation damage. Nature. 1978 Feb 16;271(5646):660-2. PMID: 342974.

11. Sun J, Chen Y, Li M, Ge Z. Role of antioxidant enzymes on ionizing radiation resistance. Free Radic Biol Med. 1998 Mar 1;24(4):586-93. PMID: 9559871.

12. Pauling L. Evolution and the need for ascorbic acid. Proc Natl Acad Sci U S A. 1970 Dec;67(4):1643-8. PMID: 5275366.

13. Natural Standard Database. Astaxanthin. http://www.naturalstandard.com/databases/herbssupplements/as-taxanthin.asp?#undefined. Accessed November 6, 2012.

14. Zhao W, Jing X, Chen C, Cui J, Yang M, Zhang Z. [Protective effects of astaxanthin against oxidative damage induced by 60Co gamma-ray irradiation]. Wei Sheng Yan Jiu. 2011 Sep;40(5):551-4. Chinese. PMID: 22043699.

15. Hofer M, Pospíšil M. Modulation of animal and human hematopoiesis by β-glucans: a review. Molecules. 2011 Sep 15;16(9):7969-79. PMID: 21921869.

16. Qi C, Cai Y, Gunn L, et al. Differential pathways regulating innate and adaptive antitumor immune responses by particulate and soluble yeast-derived ß-glucans. Blood. 2011 Jun 23;117(25):6825-36. PMID: 21531981.

17. Petruczenko A. Glucan effect on the survival of mice after radiation exposure. Acta Physiol Pol. 1984 May-Jun;35(3):231-6. PMID: 6537716.

18. Chertkov KS, Davydova SA, Nesterova TA, Zviagintseva TN, Eliakova LA. [Efficiency of polysaccharide translam for early treatment of acute radiation illness]. Radiats Biol Radioecol. 1999 Sep-Oct; 39(5):572-7. Russian. PMID: 10576030.

19. Gu YH, Takagi Y, Nakamura T, et al. Enhancement of radioprotection and anti-tumor immunity by yeast-derived beta-glucan in mice. J Med Food. 2005 Summer;8(2):154-8. PMID: 16117606.

20. Rondanelli M, Opizzi A, Monteferrario F. [The biological activity of beta-glucans]. Minerva Med. 2009 Jun;100(3):237-45. Review. Italian. PMID: 19571787.

21. Gekeler W, Grill E, Winnacker EL, et al. Algae sequester heavy metals via synthesis of phytochelatin complexes. Archives of Microbiology. 1988;150(2): 197-202.

22. Mohd Azamai ES, Sulaiman S, Mohd Habib SH, et al. Chlorella vulgaris triggers apoptosis in hepatocarcinogenesis-induced rats. J Zhejiang Univ Sci B. 2009 Jan;10(1):14-21. PMID: 19198018.

23. Makpol S, Yaacob N, Zainuddin A, Yusof YA, Ngah WZ. Chlorella vulgaris modulates hydrogen peroxide-induced DNA damage and telomere shortening of human fibroblasts derived from different aged individuals. Afr J Tradit Complement Altern Med. 2009 Jul 3;6(4):560-72. PMID: 20606778.

24. Takekoshi H, Suzuki G, Chubachi H, Nakano M. Effect of Chlorella pyrenoidosa on fecal excretion and liver accumulation of polychlorinated dibenzo-p-dioxin in mice. Chemosphere. 2005 Apr;59(2):297-304. PMID:15722102.

25. Li L, Li W, Kim YH, Lee YW. Chlorella vulgaris extract ameliorates carbon tetrachloride-induced acute hepatic

injury in mice. Exp Toxicol Pathol. 2011 Jul 7. [Epub ahead of print] PMID: 21741806. 26. Shim JA, Son YA, Park JM, Kim MK. Effect of Chlorella intake on Cadmium metabolism in rats. Nutr Res Pract. 2009 Spring;3(1):15-22. PMID: 20016697.

27. Mercola J, Klinghardt D. Mercury Toxicity and Systemic Elimination Agents. Journal of Nutritional & Environmental Medicine. 2001. 11:53-62.

28. Moison RM, Beijersbergen Van Henegouwen GM. Dietary eicosapentaenoic acid prevents systemic immunosuppression in mice induced by UVB radiation. Radiat Res. 2001 Jul;156(1):36-44. PMID: 11418071.

29. Rhodes LE, Shahbakhti H, Azurdia RM, et al. Effect of eicosapentaenoic acid, an omega-3 polyunsaturated fatty acid, on UVR-related cancer risk in humans. An assessment of early genotoxic markers. Carcinogenesis. 2003 May;24(5):919-25. PMID: 12771037.

30. Christofidou-Solomidou M, Tyagi S, Tan KS, et al. Dietary flaxseed administered post thoracic radiation treatment improves survival and mitigates radiation-induced pneumonopathy in mice. BMC Cancer. 2011 Jun 24;11:269. PMID: 21702963.

31. Weiss JF, Landauer MR. Protection against ionizing radiation by antioxidant nutrients and phytochemicals. Toxicology. 2003 Jul 15;189(1-2):1-20. Review. PMID: 12821279.

32. Mahmood J, Jelveh S, Calveley V, Zaidi A, Doctrow SR, Hill RP. Mitigation of Lung Injury after Accidental Exposure to Radiation. Radiat Res. 2011 Oct 20. PMID: 22013884.

33. El-Missiry MA, Fayed TA, El-Sawy MR, El-Sayed AA. Ameliorative effect of melatonin against gamma-irradiation-induced oxidative stress and tissue injury. Ecotoxicol Environ Saf. 2007 Feb;66(2):278-86. PMID:16793135.

34. Dreher F, Gabard B, Schwindt DA, Maibach HI. Topical melatonin in combination with vitamins E and C protects skin from ultraviolet-induced erythema: a human study in vivo. Br J Dermatol. 1998 Aug;139(2):332-9. PMID: 9767255.

35. Krinsky DL, LaValle JB, Hawkins EB, Pelton R, Ashbrook Willis N. Natural Therapeutics Pocket Guide. 2nd Edition. Hudson, Ohio: Lexi-Comp, In.c 2003.

36. Waldron-Edward D. Skoryna SC, Paul TM. Studies on the Inhibition of Intestinal Absorption of Radioactive Strontium. 3. The Effect of Administration of Sodium Alginate in Food and in Drinking Water. Can Med Assoc J. 1964 Nov 7;91:1006-10. PMID: 14222668.

37. Paul TM, Edward DW, Skoryna SC. Studies on Inhibition of Intestinal Absorption of Radioactive Strontium. II. Effects of Administration of Sodium Alginate by Orogastric Intubation and Feeding. Can Med Assoc J. 1964 Sep 5;91:553-7. PMID: 14176062.

38. Qishen P, Guo BJ, Kolman A. Radioprotective effect of extract from Spirulina platensis in mouse bone marrow cells studied by using the micronucleus test. Toxicol Lett. 1989 Aug;48(2):165-9. PMID: 2505406.

39. Mercola. Spirulina. http://articles.mercola.com/sites/articles/archive/2011/11/09/spirulina-reversed-radia-tion-damage-in-chernobyl-children-in-just-20-days.aspx. Accessed November 6, 2011.

40. Loseva, LP, Dardynskaya IV. 1993. Spirulina natural sorbent of radionucleides. Research Institute of Radiation Medicine, Minsk, Belarus. Paper presented at the 6th International Congress of Applied Algology, Czech Republic.

41. Loseva, LP. 1999. Spirulina platensis and specialties to support detoxifying pollutants and to strengthen the immune system. Paper presented at the 8th International Congress on Applied Algology. Italy.

42. Dartsch PC. Antioxidant potential of selected Spirulina platensis preparations. Phytother Res. 2008 May;22(5):627-33. PMID: 18398928.

43. Hayes DP. The protection afforded by vitamin D against low radiation damage. Int. J. Low Radiation. Vol. 5, No. 4, 2008

44. Garland CF, French CB, Baggerly LL, Heaney RP. Vitamin D supplement doses and serum 25-hydroxyvitamin D in the range associated with cancer prevention. Anticancer Res. 2011 Feb;31(2):607-11. PMID: 21378345.

45. Mizik P, Hrusovský J, Tokosová M. [The effect of natural zeolite on the excretion and distribution of radiocesium in rats]. Vet Med (Praha). 1989 Aug;34(8):467-74. Slovak. PMID: 2552638.

46. Sato I, Kudo H, Tsuda S. Removal efficiency of water purifier and adsorbent for iodine, cesium, strontium, barium and zirconium in drinking water. J Toxicol Sci. 2011;36(6):829-34. PMID: 22129747.

ABOUT THE AUTHORS





Beth Ellen DiLuglio, MS, RDN, CCN, LDN is a Certified Clinical Nutritionist (CCN) through the International and American Associations of Clinical Nutritionists (IAACN), as well as a Registered Dietitian Nutritionist (RDN) with a 20-year history of certification in nutrition support. Beth earned a Master of Science degree in Human Nutrition from Columbia University's College of Physicians & Surgeons, taught nutrition at the college level for over a decade, and completed The Institute for Functional Medicine's "Applying Functional Medicine in Clinical Practice."

Beth worked closely with researchers for the Radiation and Public Health Project (Radiation. org) and co-authored the Radiation Exposure/ Electromagnetic Hypersensitivity chapter in Dr. Ingrid Kohlstadt's medical textbook Advancing Medicine with Food and Nutrients, 2nd ed.

Join Beth's Nutrition Mission™ JoinNutritionMission@gmail.com NutritionMission.org (850) 296-8874 Nutrition Is Your Best Health Insurance!®

Julie M. Brown, MA, LMHC, is a licensed psychotherapist with a Master's degree in Counseling Psychology. She specializes in Stress Management, Women's Issues and Mind-Body Medicine and is certified in Advanced EMDR (Eye Movement Desensitization and Reprogramming) and Hypnosis.

For over two decades, Julie has focused her mind-body work on assisting cancer patients to empower and heal themselves, incorporating a variety of integrative healing modalities such as diet, somatic release, breathwork, meditation, visualization and EMDR. She is currently developing a comprehensive Cancer Prevention Resource Guide, which will bring together evidence-based tools and practices and make them available to the public via the Internet.

She served as Resource Director for the national baby teeth study, providing educational resources to parents of children with cancer. She lives in Santa Barbara, California, where she is a life coach.

Phone: (305) 321-7016 Email: juliemarlane@gmail.com

